

Part 6—References and SEBSCC Publications*

- Alexander, V., and H.J. Niebauer (1981): Oceanography at the eastern Bering Sea ice edge zone in spring. *Limnol. Oceanogr.*, *26*, 1111–1125.
- Anderson, P.J., and J.F. Piatt (1999): Community reorganization in the Gulf of Alaska following ocean climate regime shift. *Mar. Ecol. Prog. Ser.*, *189*, 117–123.
- Antonelis, G.A., E.H. Sinclair, R.R. Ream, and B.W. Robson (1997): Inter-island variation in the diet of female northern fur seals (*Callorhinus ursinus*) in the Bering Sea. *J. Zool. Soc. Lond.*, *242*, 435–451.
- Baduini, C.L., K.D. Hyrenbach, K.O. Coyle, A. Pinchuk, V. Mendenhall, and G.L. Hunt, Jr. (2001a): Mass mortality of short-tailed shearwaters in the eastern Bering Sea during summer 1997. *Fish. Oceanogr.*, *10*, 117–130.
- Baduini, C.L., J.R. Lovvorn, and G.L. Hunt, Jr. (2001b): Determining the body condition of short-tailed shearwaters: Implications for migratory flight ranges and starvation events. *Mar. Ecol. Prog. Ser.*, *222*, 265–277.
- * Baier, C.T., and J.M. Napp (2003): Climate-induced variability in *Calanus marshallae* populations. *J. Plankton Res.*, *25*, 771–782.
- Bailey, K.M., A.L. Brown, M.M. Yoklavich, and K.L. Mier (1996): Interannual variability of growth of larval and juvenile walleye pollock in the western Gulf of Alaska. *Fish. Oceanogr.*, *5*(suppl. 1), 137–147.
- Bailey, K.M., M.F. Canino, J.M. Napp, S.M. Spring, and A.L. Brown (1995): Contrasting years of prey levels, feeding condition and mortality of larval walleye pollock (*Theragra chalcogramma*) in the western Gulf of Alaska. *Mar. Ecol. Prog. Ser.*, *119*, 11–13.
- Bailey, E.P., and G.H. Davenport (1972): Die-off of murres on the Alaska Peninsula and Unimak Island. *Condor*, *74*, 215–219.
- Bailey, K.M., and S.A. Macklin (1994): Analysis of patterns in larval walleye pollock *Theragra chalcogramma* and wind mixing events in Shelikof Strait, Alaska. *Mar. Ecol. Prog. Ser.*, *113*, 1–12.
- Balch, W.M., P.M. Holligan, S.G. Ackleson, and K.J. Voss (1991): Biological and optical properties of mesoscale coccolithophore blooms in the Gulf of Maine. *Limnol. Oceanogr.*, *36*, 629–643.
- Balch, W.M., P.M. Holligan, and K.A. Kilpatrick (1992): Calcification, photosynthesis and growth of the bloom-forming coccolithophore, *Emiliania huxleyi*. *Cont. Shelf Res.*, *12*, 1353–1374.
- Balykin, P.A. (1996): Dynamics and abundance of western Bering Sea walleye pollock. In *Ecology of the Bering Sea: A Review of Russian Literature*, O.A. Mathisen, and K.O. Coyle (eds.), University of Alaska Sea Grant Report 96-01, Fairbanks, Alaska, 177–182.
- Baretta, L., and G.L. Hunt, Jr. (1994): Changes in the numbers of cetaceans near the Pribilof Islands, Bering Sea, between 1975–78 and 1987–89. *Arctic*, *47*, 321–326.
- Barnston, A.G., and R.E. Livezey (1987): Classification, seasonality and persistence of low-frequency atmospheric circulation patterns. *Mon. Weather Rev.*, *115*, 1083–1126.
- Beamish, R.J. (1995): Climate change and northern fish populations. *Can. Spec. Publ. Fish. Aquat. Sci.*, *121*, 739.
- Beamish, R.J., K.D. Leask, O.A. Ivanov, A.A. Balanov, A.M. Orlov, and E.H. Sinclair (1999a): The ecology, distribution, and abundance of midwater fishes of the Subarctic Pacific gyres. *Prog. Oceanogr.*, *43*, 399–442.

*SEBSCC publication

- Beamish, R.J., D.J. Noakes, G.A. McFarlane, L. Klyashtorin, V.V. Ivanov, and V. Kurashov (1999b): The regime concept and natural trends in the production of Pacific salmon. *Can. J. Fish. Aquat. Sci.*, 56, 516–526.
- * Blood, D.M. (2002): Low-temperature incubation of walleye pollock (*Theragra chalcogramma*) eggs from the southeast Bering Sea shelf and Shelikof Strait, Gulf of Alaska. *Deep-Sea Res. II, Top. Stud. Oceanogr.*, 49(26), 6095–6108.
- * Bond, N.A., and J.M. Adams (2002): Atmospheric forcing of the southeast Bering Sea shelf during 1995–99 in the context of a 40-year historical record. *Deep-Sea Res. II, Top. Stud. Oceanogr.*, 49(26), 5869–5887.
- * Bond, N.A., and J.E. Overland (2004): The importance of episodic weather events to the ecosystem of the Bering Sea shelf. *Fish. Oceanogr.* (in press).
- Braun, B.M., and G.L. Hunt (1983): Brood reduction in black-legged kittiwakes. *Auk*, 100, 469–476.
- * Brodeur, R.D. (2001): Habitat-specific distributions of Pacific ocean perch (*Sebastes alutus*) in Pribilof Canyon, Bering Sea. *Cont. Shelf Res.*, 21(3), 207–224.
- * Brodeur, R.D., and K.M. Bailey (2001): A different approach to understanding causes of recruitment variability in marine fishes. In *Fisheries Oceanography: An Integrative Approach to Fisheries Ecology and Management*, P.J. Harrison, and T.R. Parsons (eds.), Fish and Aquatic Resources, Series 4, Blackwell Science, 24–28.
- * Brodeur, R.D., C.E. Mills, J.E. Overland, G.E. Walters, and J.D. Schumacher (1999a): Evidence for a substantial increase in gelatinous zooplankton in the Bering Sea, with possible links to climate change. *Fish. Oceanogr.*, 8(4), 296–306.
- * Brodeur, R.D., H. Sugisaki, and G.L. Hunt, Jr. (2002): Increases in jellyfish biomass in the Bering Sea: implications for the ecosystem. *Mar. Ecol. Prog. Ser.*, 233, 89–103.
- Brodeur, R.D., and M.T. Wilson (1999): Pre-recruit walleye pollock in the eastern Bering Sea and Gulf of Alaska ecosystems. In *Proceedings of GLOBEC International Marine Science Symposium on Ecosystem Dynamics*, 238–251.
- Brodeur, R.D., P.A. Livingston, T.R. Loughlin, and A.B. Hollowed (eds.) (1996): Ecology of juvenile walleye pollock, *Theragra chalcogramma*. NOAA Tech. Rept. NMFS-126, 227 pp.
- * Brodeur, R.D., M.T. Wilson, L. Ciannelli, M.J. Doyle, and J.M. Napp (2002): Interannual and regional variability in distribution and ecology of juvenile pollock and their prey in frontal structures of the Bering Sea. *Deep-Sea Res. II, Top. Stud. Oceanogr.*, 49(26), 6051–6067.
- Brodeur, R.D., M.T. Wilson, G.E. Walters, and I.V. Melnikov (1999b): Forage fish in the Bering Sea: Distribution, species associations and biomass trends. In *Dynamics of the Bering Sea*, T.R. Loughlin, and K. Ohtani (eds.), University of Alaska Sea Grant, AK-SG-99-03, Fairbanks, Alaska, 509–536.
- Brown, C.W., and G.P. Podesta (1997): Remote sensing of coccolithophore blooms in the western South Atlantic Ocean. *Remote Sens. Environ.*, 83–91.
- Brown, C.W., and J.A. Yoder (1993): Blooms of *Emiliania huxleyi* (Prymnesiophyceae) in surface waters of the Nova Scotian shelf and the Grand Bank. *J. Plankton Res.*, 15, 1429–1438.
- Bruegeman, J.J., G.W. Oliver, C. Fowler, K. Frost, K. Fay, C. Cowles, K. Schneider, and D. Calkins (1987): Marine Mammals. In *The St. George Basin Environment and Possible Consequences of Planned Offshore Oil and Gas Development*, M.J. Hameedi (ed.), U.S. Depart. Commerce, U.S. Dept. Interior, Outer Continental Shelf Environmental Assessment Program, Juneau, Alaska, 55–81.

- Buckley, T.W., A. Greig, J. Ianelli, P. Livingston, and G. Walters (2001): Spatial distribution and ontogenetic movement of walleye pollock in the eastern Bering Sea. Presented at the North Pacific Marine Science Organization (PICES), Victoria, B.C., Canada, October 5–13, 2001, Alaska Fisheries Science Center, 7600 Sandpoint Way NE, Seattle, WA 98115.
- Bulatov, O.A. (1995): Biomass variations of walleye pollock in the Bering Sea in relation to oceanological conditions. In *Climate Change and Northern Fish Populations*, R.J. Beamish (ed.), *Can. Spec. Publ. Fish. Aquat. Sci.*, 121, 631–640.
- * Busby, M.S., K.L. Mier, and R.D. Brodeur. Habitat associations of demersal fishes and crabs in the Bering Sea near the Pribilof Islands, Alaska. *Fish. Bull.* (submitted).
- Capriulo, G.M., E.B. Sherr, and B.F. Sherr (1991): Trophic behavior and related community feeding activities of heterotrophic marine protists. In *Protozoa and Their Role in Marine Processes*, P.C. Reid, C.M. Turley, and P.H. Burkil (eds.), Springer, Berlin, 219–265.
- Cavalieri, D.J., and S. Martin (1994): The contribution of Alaskan, Siberian and Canadian coastal polynyas to the cold halocline layer of the Arctic Ocean. *J. Geophys. Res.*, 99, 18,343–18,362.
- Ciannelli, L. (2002): Cross-scale analysis of the Pribilof Archipelago, southeast Bering Sea, with a focus on age-0 walleye pollock (*Theragra chalcogramma*). Ph.D. Dissertation, University of Washington, Seattle, WA 98195, 226 pp.
- * Ciannelli, L. (2002): Effects of spatial variability, associated with a frontal structure, on predictions of age-0 walleye pollock (*Theragra chalcogramma*) growth around the Pribilof Islands, Bering Sea. *Estuar. Coast. Shelf Sci.*, 55(1), 151–165.
- * Ciannelli, L., R.D. Brodeur, and J.M. Napp (2004): Foraging impact on zooplankton by age-0 walleye pollock (*Theragra chalcogramma*) around a front in the southeast Bering Sea. *Mar. Biol.*, 144, 515–526.
- * Ciannelli, L., R.D. Brodeur, G.L. Swartzman, and S.A. Salo (2002): Physical and biological factors influencing the spatial distribution of age-0 walleye pollock (*Theragra chalcogramma*) around the Pribilof Islands, Bering Sea. *Deep-Sea Res. II, Top. Stud. Oceanogr.*, 49(26), 6109–6126.
- * Ciannelli, L., A.J. Paul, and R.D. Brodeur (2002): Regional, interannual and size-related variation of age 0 year walleye pollock whole body energy content around the Pribilof Islands, Bering Sea. *J. Fish Biol.*, 60(5), 1267–1279.
- * Ciannelli, L., B.W. Robson, R.C. Francis, K. Aydin, and R.D. Brodeur (2004): Boundaries of open marine ecosystems: An application to the Pribilof Archipelago, southeast Bering Sea ecological applications. *Ecol. Appl.*, 14(3), 942–953.
- Coachman, L.K. (1986): Circulation, water masses and fluxes on the southeastern Bering Sea shelf. *Cont. Shelf Res.*, 5, 23–108.
- Coachman, L.K., K. Aagaard, and R.B. Tripp (1975): Bering Strait: Regional physical oceanography. University of Washington Press, Seattle, WA, 172 pp.
- * Coachman, L.K., T.E. Whitledge, and J.J. Goering (1999): Silica in Bering Sea deep and bottom water. In *Dynamics of the Bering Sea*, T.R. Loughlin, and K. Ohtani (eds.), University of Alaska Sea Grant, Fairbanks, Alaska, AK-SG-99-03, 285–310.
- Conners, M.E., A.B. Hollowed, and E. Brown (2002): Retrospective analysis of Bering Sea bottom trawl surveys: Regime shift and ecosystem reorganization. *Prog. Oceanogr.*, 55, 209–222.
- Cooney, R.T., J.R. Allen, M.A. Bishop, D.L. Eslingler, T. Kline, B.L. Norcross, C.P. McRoy, J. Milton, J. Olsen, V. Patrick, A.J. Paul, D. Salmon, D. Schell, G.L. Thomas, S.L. Vaughan, and T.M. Willette (2001): Ecosystem controls of

- juvenile pink salmon (*Onchorynchus gorbuscha*) and Pacific herring (*Clupea pallasi*) populations in Prince William Sound, Alaska. *Fish. Oceanogr.*, 10(suppl. 1), 1–13.
- Cooney, R.T., and K.O. Coyle (1982): Trophic implications of cross-shelf copepod distributions in the southeastern Bering Sea. *Mar. Biol.*, 70, 187–196.
- Corkett, C.J., and I.A. McLaren (1978): The biology of *Pseudocalanus*. *Adv. Mar. Biol.*, 15, 1–231.
- * Coyle, K.O., and A.I. Pinchuk (2002): The abundance and distribution of euphausiids and zero-age pollock on the inner shelf of the southeast Bering Sea near the inner front in 1997–1999. *Deep-Sea Res. II, Top. Stud. Oceanogr.*, 49(26), 6009–6030.
- * Coyle, K.O., and A.I. Pinchuk (2002): Climate-related differences in zooplankton density and growth on the inner shelf of the southeastern Bering Sea. *Prog. Oceanogr.*, 55(1–2), 177–194.
- Cullen, J.T., Y. Rosenthal, and P.G. Falkowski (2001): The effect of anthropogenic CO₂ on the carbon isotope composition of marine phytoplankton. *Limnol. Oceanogr.*, 46, 996–998.
- Dagg, M.J., M.E. Clarke, T. Nishiyama, and S.L. Smith (1984): Production and standing stock of copepod nauplii, food items for larvae of the walleye pollock *Theragra chalcogramma* in the southeastern Bering Sea. *Mar. Ecol. Prog. Ser.*, 19, 7–16.
- * Davis, R.F., G. Lazin, J. Bartlett, A.M. Ciotti, and P.J. Stabeno (1997): Remote sensing of a pigment patch in the southeastern Bering Sea. In *Ocean Optics XIII*, S.G. Ackleson (ed.), SPIE, International Society for Optical Engineering, Vol. 2963, 654–657.
- Decker, M.B., G.L. Hunt, Jr., and G.V. Byrd, Jr. (1995): The relationships among sea surface temperature, the abundance of juvenile walleye pollock (*Theragra chalcogramma*), and the reproductive performance and diets of seabirds at the Pribilof Islands, southeastern Bering Sea. In *Climate Change and Northern Fish Populations*, R.J. Beamish (ed.), *Can. Spec. Publ. Fish. Aquat. Sci.*, 121, 425–437.
- Dragoo, D.E., G.V. Byrd, and D.B. Irons (2001): Breeding status, population trends and diets of seabirds in Alaska, 2000. U.S. Fish and Wildlife Service Report AMNWR 2001/07, 77 pp.
- * Duffy-Anderson, J.T., M.S. Busby, K.L. Mier, and C. Deliyanides. Spatial and temporal patterns in summer ichthyoplankton assemblages on the eastern Bering Sea shelf from 1996–2000. *Fish. Oceanogr.* (submitted).
- Dussart, B.M. (1965): Les différentes catégories de plancton. *Hydrobiologica*, 26, 72–74.
- Dwyer, D.A., K.M. Bailey, and P.A. Livingston (1987): Feeding habits and daily ration of walleye pollock (*Theragra chalcogramma*) in the eastern Bering Sea, with special reference to cannibalism. *Can. J. Fish. Aquat. Sci.*, 44, 1972–1984.
- Ebbesmeyer, C.C., D.R. Cayan, D.R. Milan, F.H. Nichols, D.H. Peterson, and K.T. Redmond (1991): 1976 step in the Pacific climate: Forty environmental changes between 1968–1975 and 1977–1984. In *Proceedings of the Seventh Annual Climate (PACLIM) Workshop, April 1990*, J.L. Betancourt, and V.L. Tharp (eds.), California Department of Water Resources, Interagency Ecological Studies Program Technical Report 26, 115–126.
- Eslinger, D.L., and R.L. Iverson (2001): The effects of convective and wind-driven mixing on spring phytoplankton dynamics in the southeastern Bering Sea middle shelf domain. *Cont. Shelf Res.*, 21, 627–650.

- * Flint, M.V., I.N. Sukhanova, A.I. Kopylov, S.G. Poyarkov, and T.E. Whittlestone (2002): Plankton distribution associated with frontal zones in the vicinity of the Pribilof Islands. *Deep-Sea Res. II, Top. Stud. Oceanogr.*, 49(26), 6069–6093.
- Foukal, P. (2003): Can slow variations in solar luminosity provide the missing link between the sun and climate? *Eos Trans. AGU*, 84(22), 205, 208.
- Francis, R.C., and K.M. Bailey (1983): Factors affecting recruitment of selected gadoids in the northeastern Pacific and east Bering Sea. In *Year to Year: Interannual Variability of the Environment and Fisheries of the Gulf of Alaska and the Eastern Bering Sea*, W.S. Wooster (ed.), Washington Sea Grant, University of Washington, Seattle, WA, 17–22.
- Francis, R.C., S.R. Hare, A.B. Hollowed, and W.S. Wooster (1998): Effects of interdecadal climate variability on the oceanic ecosystems of the NE Pacific. *Fish. Oceanogr.*, 7, 1–21.
- Goebel, M.E. (2002): Northern fur seal lactation, attendance and reproductive success in two years of contrasting oceanography. Ph.D. Thesis, University of California, Santa Cruz, 213 pp.
- Golet, G.H., and D.B. Irons (1999): Raising young reduces body condition and fat stores in black-legged kittiwakes. *Oecologia*, 120, 530–538.
- Golet, G.H., D.B. Irons, and J.A. Estes (1998): Survival costs of chick rearing in black-legged kittiwakes. *J. Anim. Ecol.*, 67, 827–841.
- Gower, J.R.F. (1997): Bright plankton blooms on the west coast of North America observed with AVHRR imagery. In *Monitoring Algal Blooms: New Techniques for Detecting Large-Scale Environmental Change*, M. Kahru, and C.W. Brown (eds.), Landes Bioscience, Austin, Texas, 25–41,
- Grebmeier, J.M. (1992): Benthic processes on the shallow continental shelf of the northern Bering and Chukchi Seas. In *Results of the Third Joint US-USSR Bering and Chukchi Seas Expedition (BERPAC), Summer 1988*, P.A. Nagel (ed.), U.S. Fish and Wildlife Service, Washington, DC, 243–251.
- Grebmeier, J.M., and L.W. Cooper (1994): A decade of benthic research on the continental shelves of the northern Bering and Chukchi Seas: Lessons learned. In *Bridges of Science Between North America and the Russian Far East*, R.H. Meehan, V. Serienko, and G. Weller (eds.), Fairbanks, Alaska: American Association for the Advancement of Science, Arctic Division, 87–98.
- Grebmeier, J.M., and L.W. Cooper (2002): Benthic processes in the Bering Strait region of the Arctic: Temporal/spatial variability and global change. *Eos Trans. AGU*, 83(4), Ocean Sciences Meeting Supplement, Abstract OS32M-01.
- Grebmeier, J.M., and K.H. Dunton (2000): Benthic processes in the northern Bering/Chukchi Seas: Status and global change. In *Impacts of Changes in Sea Ice and Other Environmental Parameters in the Arctic*, H.P. Huntington (ed.), Report of the Marine Mammal Commission Workshop, 15–17 February 2000, Girdwood Alaska. Available from the Marine Mammal Commission, Bethesda, Maryland.
- Haidvogel, D.B., H.G. Arango, K. Hedstrom, A. Beckmann, P. Malanotte-Rizzoli, and A.F. Shchepetkin (2000): Model evaluation experiments in the North Atlantic basin: Simulations in nonlinear terrain-following coordinates. *Dyn. Atmos. Oceans*, 32, 239–281.
- Hare, S.R., and R.C. Francis (1995): Climate change and salmon production in the northeast Pacific Ocean. In *Climate Change and Northern Fish Populations*, R.J. Beamish (ed.), *Can. Spec. Publ. Fish. Aquat. Sci.*, 121, 357–372.
- Hare, S.R., and N.J. Mantua (2000): Empirical evidence for North Pacific regime shifts in 1977 and 1989. *Prog. Oceanogr.*, 47, 103–146.

- Hare, S.R., N.J. Mantua, and R.C. Francis (1999): Inverse production regimes: Alaskan and West Coast Pacific salmon. *Fisheries*, 24, 6–14.
- Hastie, T., and R. Tibshirani (1990): Generalized additive models. Chapman and Hall, London.
- Hatch, S.A. (1987): Did the 1982–1983 El Niño-Southern Oscillation affect seabirds in Alaska? *Wilson Bull.*, 99(3), 469–474.
- * Hermann, A.J., D.B. Haidvogel, D. Musgrave, and P.J. Stabeno (2002): A regional tidal/subtidal circulation model of the southeastern Bering Sea: development, sensitivity analyses and hindcasting. *Deep-Sea Res. II, Top. Stud. Oceanogr.*, 49(26), 5945–5967.
- Hermann, A.J., P.J. Stabeno, D.B. Haidvogel, and D.L. Musgrave (2002): A regional tidal/subtidal circulation model of the southeastern Bering Sea: Development, sensitivity analyses and hindcasting. *Deep-Sea Res. II, Top. Stud. Oceanogr.*, 49, 5945–5967.
- Hinckley, S. (1987): The reproductive biology of walleye pollock, *Theragra chalcogramma*, in the Bering Sea, with reference to spawning stock structure. *Fish. Bull.*, 85, 481–498.
- Hirons, A.C., D.M. Schell, and B.P. Finny (2001): Temporal records of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ in North Pacific pinnipeds: Inferences regarding environmental change and diet. *Oecologia*, 129, 591–601.
- Holligan, P.M., M. Viollier, D.S. Harbour, P. Camus, and M. Champagne-Philippe (1983): Satellite and ship studies of coccolithophore production along a continental shelf edge. *Nature*, 304, 339–342.
- Hollowed, A.B., S.R. Hare, and W.S. Wooster (2001): Pacific basin climate variability and patterns of northeast Pacific marine fish production. *Prog. Oceanogr.*, 49, 257–282.
- Hollowed, A.B., and W.S. Wooster (1995): Decadal-scale variations in the eastern subarctic Pacific: II Responses of Northeast Pacific fish stocks. In *Climate Change and Northern Fish Populations*, R.J. Beamish (ed.), *Can. Spec. Publ. Fish. Aquat. Sci.*, 121, 373–385.
- Honkalehto, T., W. Patton, S. De Blois, and N. Williamson (2002): Echo integration-trawl survey results for walleye pollock (*Theragra chalcogramma*) on the Bering Sea shelf and slope during summer 2000. NOAA Tech. Memo. NMFS-AFSC-126, 66 pp.
- Hood, D.W. (1999): PROBES: Processes and resources of the eastern Bering Sea shelf. In *Dynamics of the Bering Sea*, T.R. Loughlin, and K. Ohtani (eds.), University of Alaska Sea Grant, AK-SG-99-03, Fairbanks, AK, 387–407.
- Howell-Kubler, A., E. Lessard, and J. Napp (1996): Springtime microprotozoan abundance and biomass in the southeastern Bering Sea and Shelikof Strait, Alaska. *J. Plankton Res.*, 18, 731–745.
- * Hunt, G.L., Jr., C.L. Baduini, R.D. Brodeur, K.O. Coyle, N.B. Kachel, J.M. Napp, S.A. Salo, J.D. Schumacher, P.J. Stabeno, D.A. Stockwell, T.E. Whittle, and S.I. Zeeman (1999): The Bering Sea in 1998: A second consecutive year of extreme weather-forced anomalies. *Eos Trans. AGU*, 80(561), 565–566.
- * Hunt, G.L., Jr., C.L. Baduini, and J. Jahncke (2002): Diets of short-tailed shearwaters in the southeastern Bering Sea. *Deep-Sea Res. II, Top. Stud. Oceanogr.*, 49(26), 6147–6156.
- * Hunt, G.L., Jr., and G.V. Byrd, Jr. (1999): Marine bird populations and carrying capacity of the eastern Bering Sea. In *Dynamics of the Bering Sea*, T.R. Loughlin, and K. Ohtani (eds.), University of Alaska Sea Grant, Fairbanks, Alaska, AK-SG-99-03, 631–650.

- Hunt, G.L., Jr., K.O. Coyle, S. Hoffman, M.B. Decker, and E.N. Flint (1996a): Foraging ecology of short-tailed shearwaters near the Pribilof Islands, Bering Sea. *Mar. Ecol. Prog. Ser.*, 141, 1–11.
- Hunt, G.L., Jr., M.B. Decker, and A.S. Kitaysky (1996b): Fluctuations in the Bering Sea ecosystem as reflected in the reproductive ecology and diets of kittiwakes on the Pribilof Islands, 1975 to 1990. In *Aquatic Predators and Their Prey*, S. Greenstreet, and M. Tasker (eds.), Blackwell, London, 142–153.
- Hunt, G.L., Jr., and N.M. Harrison (1990): Foraging habitat and prey taken by Least Auklets at King Island, Alaska. *Mar. Ecol. Prog. Ser.*, 65, 141–150.
- Hunt, G.L., Jr., N.M. Harrison, and T. Cooney (1990): Foraging of Least Auklets: The influence of hydrographic structure and prey abundance. *Stud. Avian Biol.*, 14, 7–22.
- Hunt, G.L., Jr., A.S. Kitaysky, M.B. Decker, D.E. Dragoo, and A.M. Springer (1996c): Changes in the distribution and size of juvenile walleye pollock as indicated by seabird diets at the Pribilof Islands and by bottom trawl surveys in the eastern Bering Sea. In *Ecology of Juvenile Walleye Pollock, Theragra chalcogramma*, R.D. Brodeur, P.A. Livingston, T.R. Loughlin, and A.B. Hollowed (eds.), NOAA Tech. Rept. NMFS-126, 133–147.
- * Hunt, G.L., Jr., and P.J. Stabeno (2002): Climate change and the control of energy flow in the southeastern Bering Sea. *Prog. Oceanogr.*, 55(1–2), 5–22.
- * Hunt, G.L., Jr., P.J. Stabeno, G.E. Walters, E.H. Sinclair, R.D. Brodeur, J.M. Napp, and N.A. Bond (2002): Climate change and control of the southeastern Bering Sea pelagic ecosystem. *Deep-Sea Res. II, Top. Stud. Oceanogr.*, 49(26), 5821–5853.
- Huntley, M.E., and M.D.G. Lopez (1992): Temperature-dependent production of marine copepods: A global analysis. *Am. Nat.*, 140, 201–242.
- Ianelli, J.N., T. Buckley, T. Honkalehto, N. Williamson and G. Walters (2001): Bering Sea-Aleutian Islands Walleye Pollock Assessment for 2002. In *Stock assessment and Fishery Evaluation Report for the Groundfish Resources of the Bering Sea/Aleutian Islands Regions*, North Pacific Fishery Management Council, Anchorage, AK, Section 1:1–105.
- Ianelli, J., L.W. Fritz, T. Honkalehto, N. Williamson and G. Walters (1998): Eastern Bering Sea walleye pollock stock assessment with yield considerations for 1999. In *Stock Assessment and Fishery Evaluation Report for the Groundfish Resources of the Bering Sea/Aleutian Islands Regions*, pp. 31–111, Groundfish Plan Team for the Bering Sea-Aleutian Islands Region (ed.), North Pacific Fishery Management Council, 605 W. 4th Ave., Suite 306, Anchorage, AK 99501, 560 pp.
- IARPC (Interagency Arctic Research Planning Committee) (2001): United States Arctic Research Plan: Biennial Revision 2002–2006. Office of Polar Programs, National Science Foundation, 84 pp.
- Iida, T., S-I. Saitoh, T. Miyamura, M. Toratani, H. Fukushima, and N. Shiga (2002): Temporal and spatial variability of coccolithophore blooms in the eastern Bering Sea, 1998–2001. *Prog. Oceanogr.*, 55, 165–175.
- Ingraham, W.J., Jr., and R.K. Miyahara (1988): Ocean surface current simulations in the North Pacific Ocean and Bering Sea (OSCURS-Numerical Model). NOAA Tech. Memo. NMFS-F/NWC-130, 155 pp.
- Irons, D.B. (1992): Aspects of foraging behavior and reproductive biology of the black-legged Kittiwake. Ph.D. Dissertation, University of California, Irvine, 143 pp.
- Iverson, R.L., L.K. Coachman, R.T. Cooney, T.S. English, J.J. Goering, G.L. Hunt, Jr., M.C. Macauley, C.P. McRoy, W.S. Reeburgh, and T.E. Whittlesey (1979a):

- Ecological significance of fronts in the southeastern Bering Sea. In *Ecological Processes in Coastal and Marine Systems*, R.J. Livingston (ed.), Plenum Press, New York, 437–466.
- Iverson, S.J., K.J. Frost, and S.L.C. Lang (2002): Fat content and fatty acid composition of forage fish and invertebrates in Prince William Sound, Alaska: Factors contributing to among and within species variability. *Mar. Ecol. Prog. Ser.*, *241*, 161–181.
- Iverson, R.L., T.E. Whitledge, and J.J. Goering (1979b): Chlorophyll and nitrate fine structure in the southeastern Bering Sea shelf break front. *Nature*, *281*, 664–666.
- * Jahncke, J., K.O. Coyle, S.I. Zeeman, N.B. Kachel, and G.L. Hunt, Jr. Distribution of foraging shear waters with respect to the inner front of the southeastern Bering Sea. *Mar. Ecol. Prog. Ser.* (submitted).
- Johnson, G.C., P.J. Stabeno, and S.C. Riser (2004): The Bering Slope Current system revisited. *J. Phys. Oceanogr.*, *34*(2), 384–398.
- * Jurado-Molina, J., and P.A. Livingston (2002): Climate forcing effects on trophically linked groundfish populations: Implications for fisheries management. *Can. J. Fish. Aquatic Sci.*, *59*, 1941–1951.
- * Kachel, N.B., G.L. Hunt, Jr., S.A. Salo, J.D. Schumacher, P.J. Stabeno, and T.E. Whitledge (2002): Characteristics and variability of the inner front of the southeastern Bering Sea. *Deep-Sea Res. II, Top. Stud. Oceanogr.*, *49*(26), 5889–5909.
- Kalnay, E., M. Kanamitsu, R. Kistler, W. Collins, D. Deaven, L. Gandin, M. Iredell, S. Saha, G. White, J. Woollen, Y. Zhu, A. Leetmaa, B. Reynolds, M. Chelliah, W. Ebisuzaki, W. Higgins, J. Janowiak, K. C. Mo, C. Ropelewski, J. Wang, R. Jenne, and D. Joseph (1996): The NCEP/NCAR 40-Year Reanalysis Project. *Bull. Am. Meteorol. Soc.*, *77*, 437–472.
- Kendall, A.W. (2001): Specific gravity and vertical distribution of walleye pollock (*Theragra chalcogramma*) eggs. AFSC Processed Report 2001–01, January 2001, 45 pp.
- Kitaysky, A.S., G.L. Hunt, Jr., E.N. Flint, M.A. Rubega, and M.B. Decker (2000): Resource allocation in breeding seabirds: Responses to fluctuations in their food supply. *Mar. Ecol. Prog. Ser.*, *206*, 283–296.
- Kitaysky, A.S., E.V. Kitaiskaia, J.C. Wingfield, and J.F. Piatt (2001a): Dietary restriction causes chronic elevation of corticosterone and enhances stress response in red-legged kittiwake chicks. *J. Comp. Physiol., B*, *171*, 701–709.
- Kitaysky, A.S., E.V. Kitaysky, J.F. Piatt, and J.C. Wingfield (2003): Benefits and costs of increased levels of corticosterone in seabird chicks. *Horm. Behav.*, *43*, 140–149.
- Kitaysky, A.S., J.F. Piatt, and J.C. Wingfield (2002): Are seabirds breeding in the southeastern Bering Sea food-limited? Final Report to the North Pacific Marine Research Program, North Pacific Research Board, Anchorage, Alaska.
- Kitaysky, A.S., J.F. Piatt, J.C. Wingfield, and M. Romano (1999a): The adrenocortical stress-response of black-legged kittiwake chicks in relation to dietary restrictions. *J. Comp. Physiol., B*, *169*, 303–310.
- Kitaysky, A.S., J.C. Wingfield, and J.F. Piatt (1999b): Dynamics of food availability, body condition and physiological stress response in breeding black-legged kittiwakes. *Funct. Ecol.*, *13*, 577–584.
- Kitaysky, A.S., J.C. Wingfield, and J.F. Piatt (2001b): Corticosterone facilitates begging and affects resource allocation in the black-legged kittiwake. *Behav. Ecol.*, *12*, 619–625.
- Klyashtorin, L.B., and F.N. Rukhlov (1998): Long-term climate change and pink salmon stock fluctuations. *N. Pac. Anadromous Fish Comm. Bull.*, *1*, 464–479.

- Kruse, G.H. (1998): Salmon run failures in 1997–98: a link to anomalous conditions? *Alaska Fish. Bull.*, 5, 55–63.
- *Ladd, C., G.L. Hunt, Jr., and P.J. Stabeno: Climate, mixing, and phytoplankton on the southeastern Bering Sea shelf: A synthesis (unpublished manuscript).
- Lang, G.M., R.D. Brodeur, J.M. Napp, and R. Schabetsberger (2000): Variation in groundfish predation on juvenile walleye pollock relative to hydrographic structure near the Pribilof Islands, Alaska. *ICES J. Mar. Sci.*, 57, 265–271.
- Lang, G.M., C.W. Derrah, and P.A. Livingston (2003): Groundfish food habits and predation on commercially important prey species in the eastern Bering Sea from 1993 to 1996. U.S. Department of Commerce AFSC Processed Report 2003–2004. Alaska Fisheries Science Center, 7600 Sand Point Way NE, Seattle, WA 98115, 351 pp. preparation).
- Leatherwood, S., A.E. Bowles, and R.R. Reeves (1983): Aerial surveys of marine mammals in the southeastern Bering Sea. National Oceanic and Atmospheric Administration/Outer Continental Shelf Environmental Assessment Program, Environmental Assessment of the Alaskan Continental Shelf, Final Reports of Principal Investigators, 42, 147–473.
- Lee, S.H. (2000): Use of Beaufort Sea as feeding habitat by bowhead whales (*Balaena mysticetus*) as indicated by stable isotope ratios. MS Thesis, University of Alaska Fairbanks, Fairbanks, AK 99775, 95 pp.
- Livingston, P.A. (ed.) (1991): Groundfish food habits and predation on commercially important prey species in the eastern Bering Sea from 1984 to 1986. NOAA Tech. Memo. NMFS-F/NWC-207, 240 pp.
- Livingston, P.A. (1993): The importance of predation by groundfish, marine mammals, and birds on walleye pollock *Theragra chalcogramma* and Pacific herring *Clupea pallasi* in the eastern Bering Sea. *Mar. Ecol. Prog. Ser.*, 102, 205–215.
- Livingston, P.A., and Y. deReynier (1996): Groundfish food habits and predation on commercially important prey species in the eastern Bering Sea from 1990 to 1992. U.S. Department of Commerce AFSC Processed Report 96-04, 214 pp.
- Livingston, P.A., and J. Jurado-Molina (2000): A multispecies virtual population analysis of the eastern Bering Sea. *ICES J. Mar. Sci.*, 57, 294–299.
- Livingston, P.A., and G.M. Lang (1996): Interdecadal comparisons of walleye pollock, *Theragra chalcogramma*, cannibalism in the eastern Bering Sea. In *Ecology of Juvenile Walleye Pollock, Theragra chalcogramma*, R.D. Brodeur, P.A. Livingston, T.R. Loughlin, and A.B. Hollowed (eds.), NOAA Tech. Rept. NMFS-126, 115–124.
- Livingston, P.A., and R.D. Methot (1998): Incorporation of predation into a population assessment model of eastern Bering Sea walleye pollock. In *Fishery Stock Assessment Models*, Alaska Sea Grant College Program, AK-SG-98-01, 16 pp.
- Livingston, P.A., L.-L. Low, and R.J. Marasco (1999): Eastern Bering Sea trends. In *Large Marine Ecosystems of the Pacific Rim: Assessment, Sustainability, and Management*, K. Sherman, and Q. Tang (eds.), Blackwell Science, Inc., Malden, MA, USA, 140–162.
- Livingston, P.A., A. Ward, G.M. Lang, and M.-S. Yang (1993): Groundfish food habits and predation on commercially important prey species in the eastern Bering Sea from 1987 to 1989. NOAA Tech. Memo. NMFS-AFSC-11, 192 pp.
- Loughlin, T.R., G.A. Antonelis, J.D. Baker, A.E. York, C.W. Fowler, R.L. DeLong, and H.W. Braham (1994): Status of the northern fur seal population in the United States during 1992. In *Fur Seal Investigations, 1994*, E.H. Sinclair (ed.), NOAA Tech. Memo. NMFS-AFSC-45.
- Mackas, D.L., R. Goldblatt, and A.G. Lewis (1998): Interdecadal variation in de-

- velopmental timing of *Neocalanus plumchrus* populations at Ocean Station P in the subarctic North Pacific. *Can. J. Fish. Aquatic Sci.*, 55, 1878–1893.
- MacKenzie, B.R., T.J. Miller, S. Cyr, and W.C. Leggett (1994): Evidence for a dome-shaped relationship between turbulence and larval fish ingestion rates. *Limnol. Oceanogr.*, 39(8), 1790–1799.
- Macklin, S.A. (1999a): Bering Sea FOCI. In *Dynamics of the Bering Sea*, T.R. Loughlin, and K. Ohtani (eds.), Univ. of Alaska Sea Grant AK-SG-99-03, Fairbanks, Alaska, 733–751.
- * Macklin, S.A. (1999b): Southeast Bering Sea Carrying Capacity (SEBSCC). *PICES Press*, 7(2), 24–28, 34.
- * Macklin, S.A. (2001): Bering Sea and North Pacific Ocean theme page: Providing Internet-based information to scientists, managers and educators for six years. *PICES Press*, 9(1), 29–30, 34.
- * Macklin, S.A., G.L. Hunt, Jr., and J.E. Overland (2002): Collaborative research on the pelagic ecosystem of the southeastern Bering Sea shelf. *Deep-Sea Res. II, Top. Stud. Oceanogr.*, 49(26), 5813–5819.
- * Macklin, S.A., V.I. Radchenko, S. Saitoh, and P.J. Stabeno (2002): Variability in the Bering Sea ecosystem. *Prog. Oceanogr.*, 55(1–2), 1–4.
- * Macklin, S.A., and P.J. Stabeno (2001): Marine ecosystem studies of physical and biological interactions in the eastern Bering Sea and western Gulf of Alaska. In *Arctic Research of the United States*, 14(2), National Science Foundation, Arlington, Virginia, 25–32.
- Mantua, N.J., and S.R. Hare (2002): The Pacific Decadal Oscillation. *J. Oceanogr.*, 58, 35–44.
- Mantua, N.J., S.R. Hare, Y. Zhang, J.M. Wallace, and R.C. Francis (1997): A Pacific interdecadal climate oscillation with impacts on salmon production. *Bull. Am. Meteorol. Soc.*, 78, 1069–1079.
- Marchesiello, P., J.C. McWilliams, and A.F. Shchepetkin (2001): Open boundary conditions for long-term integration of regional oceanic models. *Ocean Model.*, 3, 1–20.
- Martin, J.E., R.D. Grauman, and N. Marsili (2001): Surface cycloclisis in the North Pacific Ocean. Part I: A synoptic climatology. *Mon. Weather Rev.*, 129, 748–765.
- McFarlane, G.A., J.R. King, and R.J. Beamish (2000): Have there been recent changes in climate? Ask the fish. *Prog. Oceanogr.*, 47, 147–169.
- McGowan, J.A. (1990): Climate and change in oceanic ecosystems: the value of time-series data. *Trends in Ecological Evolution*, 5, 293–299.
- McLaren, I.A. (1963): Effects of temperature on growth of zooplankton and the adaptive significance of vertical migration. *J. Fish. Res. Board Can.*, 20, 685–727.
- McRoy, C.P., and J.J. Goering (1974): The influence of ice on the primary productivity of the Bering Sea. In *Oceanography of the Bering Sea With Emphasis on Renewable Resources*, D.W. Hood, and E.J. Kelly (eds.), Occasional Publication No. 2, Institute of Marine Sciences, University of Alaska, Fairbanks, Alaska, 403–421.
- Megrey, B.A., and S. Hinckley (2001): Effect of turbulence on feeding of larval fishes: a sensitivity analysis using an individual-based model. *ICES J. Mar. Sci.*, 58, 1015–1029.
- Megrey, B.A., A.B. Hollowed, S.R. Hare, S.A. Macklin, and P.J. Stabeno (1996): Contribution of FOCI research to forecasts of year-class strength of walleye pollock in Shelikof Strait, Alaska. *Fish. Oceanogr.*, 5(suppl. 1), 189–203.

*SEBSCC publication

- Merrick, R.L. (1997): Current and historical roles of apex predators in the Bering Sea ecosystem. *J. Northwest Atl. Fish. Sci.*, 22, 343–355.
- Merrick, R.L., M.K. Chumbley, and G.V. Byrd (1997): A potential relationship between the diet diversity of Steller sea lions (*Eumetopias jubatus*) and their population decline in Alaska. *Can. J. Fish. Aquatic Sci.*, 54, 1342–1348.
- Minobe, S. (1999): Resonance in bidecadal and pentadecadal climate oscillations over the North Pacific: Role in climate regime shifts. *Geophys. Res. Lett.*, 26, 855–858.
- Mitchell-Innes, B.A., and A. Winter (1987): Coccolithophores: A major component in mature upwelled waters off Cape Peninsula, South Africa in March, 1983. *Mar. Biol.*, 95, 25–30.
- Mock, D.W., H. Drummond, and C.H. Stinson (1990): Avian siblicide. *Am. Sci.*, 78, 438–449.
- Moore, S.E., J.M. Waite, N.A. Friday, and T. Honkalehto (2002): Cetacean distribution and relative abundance on the central-eastern and the southeastern Bering Sea shelf with reference to oceanographic domains. *Prog. Oceanogr.*, 55, 249–261.
- Moore, S.E., M. Waite, L. Mazzuca, and R.C. Hobbs (2000): Provisional estimates of Mysticete whale abundance on the central Bering Sea shelf. *J. Cetacean Res. Manage.*, 2, 227–234.
- Nakatani, T., and T. Maeda (1984): Thermal effect on the development of walleye pollock eggs and on their upward speed to the surface. *Bull. Jap. Soc. Sci. Fish.*, 50, 937–942.
- * Napp, J.M., C.T. Baier, K.O. Coyle, R.D. Brodeur, N. Shiga, and K.L. Mier (2002): Interannual and decadal variability in zooplankton communities of the southeast Bering Sea shelf. *Deep-Sea Res. II, Top. Stud. Oceanogr.*, 49(26), 5991–6008.
- * Napp, J.M., and G.L. Hunt, Jr. (2001): Anomalous conditions in the south-eastern Bering Sea 1997: Linkages among climate, weather, ocean, and biology. *Fish. Oceanogr.*, 10(1), 61–68.
- Napp, J.M., A.W. Kendall, Jr., and J.D. Schumacher (2000): A synthesis of biological and physical processes affecting the feeding environment of larval walleye pollock (*Theragra chalcogramma*) in the eastern Bering Sea. *Fish. Oceanogr.*, 9, 147–162.
- NCDC (National Climatic Data Center) (2004): Global surface temperature anomalies. <http://www.ncdc.noaa.gov/oa/climate/research/anomalies/anomalies.html>.
- Niebauer, H.J. (1983): Multiyear sea ice variability in the eastern Bering Sea: an update. *J. Geophys. Res.*, 88, 2733–2742.
- * Niebauer, H.J. (1998): Variability in Bering Sea ice cover as affected by a regime shift in the North Pacific in the period 1947–1996. *J. Geophys. Res.*, 103(C12), 27,717–27,737.
- Niebauer, H.J., V. Alexander, and R.T. Cooney (1981): Primary production at the eastern Bering Sea ice edge: The physical and biological regimes. In *The Eastern Bering Sea Shelf: Oceanography and Resources*, D.W. Hood, and J.A. Calder (eds.), Vol. 1, U.S. Government Printing Office, Washington, D.C., 763–780.
- Niebauer, H.J., V. Alexander, and S.M. Henrichs (1990): Physical and biological oceanographic interaction in the spring bloom at the Bering Sea marginal ice edge zone. *J. Geophys. Res.*, 95, 22,229–22,242.
- Niebauer, H.J., V. Alexander, and S.M. Henrichs (1995): A time-series study of the spring bloom at the Bering Sea ice edge I: Physical processes, chlorophyll and nutrient chemistry. *Cont. Shelf Res.*, 15, 1859–1878.

- * Niebauer, H.J., N.A. Bond, L.P. Yakunin, and V.V. Plotnikov (1999): An update on the climatology and sea ice of the Bering Sea. In *Dynamics of the Bering Sea*, T.R. Loughlin and K. Ohtani (eds.), University of Alaska Sea Grant, Fairbanks, Alaska, AK-SG-99-03, 29–60.
- Nishiyama, T. (1974): Energy requirements of sockeye salmon in Bristol Bay in the eastern Bering Sea. In *Oceanography of the Bering Sea With Emphasis on Renewable Resources*, D.W. Hood, and E.J. Kelly (eds.), Institute of Marine Sciences Occasional Publication 2, University of Alaska, Fairbanks, Alaska, 321–343.
- NMFS (National Marine Fisheries Service) (1993): Final conservation plan for the northern fur seal (*Callorhinus ursinus*). Prepared by the NMFS/Alaska Fisheries Science Center/National Marine Mammal Laboratory, Seattle, WA and the NMFS/Office of Protected Resources, Silver Spring, MD, 80 pp.
- NMFS (National Marine Fisheries Service) (1995): Status review of the United States Steller sea lion (*Eumetopias jubatus*) population. Prepared by the NMFS/Alaska Fisheries Science Center/National Marine Mammal Laboratory, Seattle, WA, 45 pp.
- NMFS (National Marine Fisheries Service) (2002): Fewer northern fur seal pups born in 2002. National Marine Fisheries Service, Alaska Region Press Release, 19 December 2002, <http://www.fakr.noaa.gov>.
- NMML (National Marine Mammal Laboratory) (1994): Status review of Steller sea lions (*Eumetopias jubatus*). National Marine Mammal Laboratory, National Marine Fisheries Service, NOAA, Seattle, WA, 40 pp.
- NPAFC (North Pacific Anadromous Fish Commission) (2001): Draft plan for NPAFC Bering-Aleutian International Survey (BASIS). NPAFC Doc. 579, Rev. 1, 27 pp.
- NPFMC (North Pacific Fishery Management Council) (2000): Stock Assessment and Fishery Evaluation document.
- NPFMC (North Pacific Fishery Management Council) (2001): Stock Assessment and Fishery Evaluation document.
- NPFMC (North Pacific Fishery Management Council) (2002): Stock Assessment and Fishery Evaluation document.
- NRC (National Research Council) (1996): The Bering Sea Ecosystem. National Academy Press, Washington, D.C., 324 pp.
- NRC (National Research Council) (2003): Decline of the Steller sea lion in Alaskan waters: Untangling food webs and fishing nets. National Academy Press, Washington, D.C.
- Ohtani, K., and T. Azumaya (1995): Influence of interannual changes in ocean conditions on the abundance of walleye pollock (*Theragra chalcogramma*) in the eastern Bering Sea. In *Climate Change and Northern Fish Populations*, R.J. Beamish (ed.), *Can. Spec. Publ. Fish. Aquat. Sci.*, 121, 87–95.
- Oka, N., and N. Maruyama (1986): Mass mortality of short-tailed shearwaters along the Japanese coast. *Tori*, 34, 97–104.
- Okkonen, S.R. (2001): Altimeter observations of the Bering Slope Current eddy field. *J. Geophys. Res.*, 106, 2465–2476.
- * Olson, M.B., and S.L. Strom (2002): Phytoplankton growth, microzooplankton herbivory and community structure in the southeast Bering Sea: Insight into the formation and temporal persistence of an *Emiliania huxleyi* bloom. *Deep-Sea Res. II, Top. Stud. Oceanogr.*, 49(26), 5969–5990.
- Overland, J.E., J.M. Adams, and N.A. Bond (1999): Decadal variability in the Aleutian low and its relation to high-latitude circulation. *J. Clim.*, 12, 1542–1548.

- * Overland, J.E., J.M. Adams, and H.O. Mofjeld (2001): Chaos in the North Pacific: spatial modes and temporal irregularity. *Prog. Oceanogr.*, *47*, 337–354.
- * Overland, J.E., N.A. Bond, and J.M. Adams (2001): North Pacific atmospheric and SST anomalies in 1997: Links to ENSO? *Fish. Oceanogr.*, *10*(1), 69–80.
- * Overland, J.E., N.A. Bond, and J.M. Adams (2002): The relation of surface forcing of the Bering Sea to large-scale climate patterns. *Deep-Sea Res. II, Top. Stud. Oceanogr.*, *49*(26), 5855–5868.
- Overland, J.E., and C.H. Pease (1982): Cyclonic climatology of the Bering Sea and its relation to sea ice extent. *Mon. Weather Rev.*, *110*, 5–13.
- Overland, J.E., M.C. Spillane, H.E. Hurlburt, and A.J. Wallcraft (1994): A numerical study of the circulation of the Bering Sea basin and exchange with the North Pacific Ocean. *J. Phys. Oceanogr.*, *24*, 736–758.
- * Overland, J.E., and P.J. Stabeno (2004): Is the climate of the Bering Sea warming and affecting the ecosystem? *Eos, Trans. AGU* (in press).
- Pease, C.H. (1980): Eastern Bering Sea ice processes. *Mon. Weather Rev.*, *108*, 2015–2023.
- Percival, D.B., J.E. Overland, and H.O. Mofjeld (2001): Interpretation of North Pacific variability as a short- and long-memory process. *J. Clim.*, *14*, 4545–4559.
- Peterson, R.T., and J. Fisher (1955): *Wild America*. Houghton Mifflin Co., Boston, MA, 434 pp.
- Peterson, W.T., J.E. Kister, and L.R. Feinberg (2002): The effects of the 1997–1999 El Niño-La Niña events on the hydrography and zooplankton off the Central Oregon coast. *Prog. Oceanogr.*, *54*, 381–398.
- Piatt, J.F., and P. Anderson (1996): Response of common murres to the *Exxon Valdez* oil spill and long-term changes in the Gulf of Alaska marine ecosystem. *Am. Fish. Soc. Symp.*, *18*, 720–737.
- Piatt, J.F., and T.I. van Pelt (1997): Mass-mortality of guillemots (*Uria aalge*) in the Gulf of Alaska in 1993. *Mar. Pollut. Bull.*, *34*, 656–662.
- Polovina, J.J. (1984): The ECOPATH model and its application to French Frigate Shoals. *Coral Reefs*, *3*, 1–11.
- Polovina, J.J., G.T. Mitchum, and G.T. Evans (1995): Decadal and basin-scale variations in mixed layer depth and the impact on biological production in the Central and North Pacific. *Deep-Sea Res.*, *42*, 1701–1716.
- Price, J.F., R.A. Weller, and R. Pinkel (1986): Diurnal cycling: Observations and models of the upper ocean response to diurnal heating, cooling and wind mixing. *J. Geophys. Res.*, *91*, 8411–8427.
- Quinn, T.J., and H.J. Niebauer (1995): Relation of eastern Bering Sea walleye pollock (*Theragra chalcogramma*) recruitment to environmental and oceanographic variables. In *Climate Change and Northern Fish Populations*, R.J. Beamish (ed.), *Can. Spec. Publ. Fish. Aquat. Sci.*, *121*, 497–507.
- Reed, R.K. (1995): On the variable subsurface environment of fish stocks in the Bering Sea. *Fish. Oceanogr.*, *4*, 317–323.
- Reed, R.K. (1998): Confirmation of a convoluted flow over the Southeastern Bering Sea Shelf. *Cont. Shelf Res.*, *18*, 99–103.
- Reed, R.K. (1999): A time series of temperature, salinity, and geopotential across the southeastern Bering Sea shelf, 1995–1999. NOAA Tech. Rept. ERL 455-PMEL 43 (NTIS PB2000-100564), 15 pp.
- Reed, R.K., and P.J. Stabeno (1996): On the climatological mean circulation over the eastern Bering Sea shelf. *Cont. Shelf Res.*, *16*(10), 1297–1305.

- Reed, R.K., and P.J. Stabeno (1999): The Aleutian North Slope Current. In *Dynamics of the Bering Sea*, T.R. Loughlin, and K. Ohtani (eds.), University of Alaska Sea Grant, AK-SG-99-03, 177–192.
- * Reed, R.K., and P.J. Stabeno (2002): Surface heat fluxes and subsurface heat content at a site over the southeastern Bering Sea shelf, May–July 1996. *Deep-Sea Res. II, Top. Stud. Oceanogr.*, 49(26), 5911–5917.
- Rho, T. (2000): Carbon and nitrogen uptake dynamics during 1997 and 1998: Anomalous conditions in the Bering Sea. M.S. Thesis, University of Alaska Fairbanks, 95 pp.
- * Rho, T., J.J. Goering, T.E. Whitledge, and D.A. Stockwell. Carbon and nitrogen uptakes and nutrients response to the warm conditions in the southeastern Bering Sea during 1997 and 1998. *Cont. Shelf Res.* (in revision).
- Rivkin, R.B., J.N. Putland, M.R. Anderson, and D. Deibel (1999): Microzooplankton bactivory and herbivory in the NE subarctic Pacific. *Deep-Sea Res. (2 Top. Stud. Oceanogr.)*, 46, 2579–2618.
- Roach, A.T., K. Aagaard, C.H. Pease, S.A. Salo, T. Weingartner, V. Pavlov, and M. Kulakov (1995): Direct measurements of transport and water properties through Bering Strait. *J. Geophys. Res.*, 100, 18,443–18,457.
- Robertson, J.E., C. Robinson, D.R. Turner, P. Holligan, A.J. Watson, P. Boyd, E. Fernandez, and M. Finch (1994): The impact of a coccolithophore bloom on oceanic carbon uptake in the northeast Atlantic during summer 1991. *Deep-Sea Res.*, 41, 297–314.
- Robson, B.W. (2002): The relationship between foraging areas and breeding sites of lactating northern fur seals, *Callorhinus ursinus* in the eastern Bering Sea. M.S. Thesis, University of Washington, Seattle, WA, 62 pp.
- Rosenkranz, G.E., A.V. Tyler, and G.H. Kruse (2001): Effects of water temperature and wind on year-class strength of Tanner crabs in Bristol Bay, Alaska. *Fish. Oceanogr.*, 10(1), 1–12.
- Royer, T.C. (1982): Coastal fresh water discharge in the Northeast Pacific. *J. Geophys. Res.*, 87, 2017–2021.
- Russell, R.W., N.M. Morrison, and G.L. Hunt, Jr. (1999): Foraging at a front: hydrography, zooplankton and avian planktivory in the northern Bering Sea. *Mar. Ecol. Prog. Ser.*, 182, 77–93.
- Sambrotto, R.N., and J.J. Goering (1983): Interannual variability of phytoplankton and zooplankton production on the southeast Bering Sea shelf. In *From Year to Year*, W.S. Wooster (ed.), Washington Sea Grant, Seattle, WA, 161–177.
- Sambrotto, R.N., H.J. Niebauer, J.J. Goering, and R.L. Iverson (1986): Relationships among vertical mixing, nitrate uptake and phytoplankton growth during the spring bloom in the Southeast Bering Sea middle shelf. *Cont. Shelf Res.*, 5, 161–198.
- * Schabetsberger, R.D., R.D. Brodeur, L. Ciannelli, J.M. Napp, and G.L. Swartzman (2000): Diel vertical migration and interaction of zooplankton and juvenile walleye pollock (*Theragra chalcogramma*) at a frontal region near Pribilof Islands, Bering Sea. *ICES J. Mar. Sci.*, 57(4), 1283–1295.
- Schell, D.M. (2000): Declining carrying capacity in the Bering Sea: Isotopic evidence from whale baleen. *Limnol. Oceanogr.*, 45, 459–462.
- Schell, D.M. (2001): Carbon isotope ratio variations in the Bering Sea biota: The role of anthropogenic carbon dioxide. *Limnol. Oceanogr.*, 46, 999–1000.
- Schell, D.M., B.A. Barnett, and K. Vinette (1998): Carbon and nitrogen isotope ratios in zooplankton of the Bering, Chukchi, and Beaufort Seas. *Mar. Ecol. Prog. Ser.*, 162, 11–23.

- Schnute, J.T. (1994): A general framework for developing sequential fisheries models. *Can. J. Fish. Aquat. Sci.*, 51, 1676–1688.
- * Schumacher, J.D. (2000): Regime shift theory: A review of changing environmental conditions in the Bering Sea and eastern North Pacific. In *Proceedings of the Fifth North Pacific Rim Fisheries Conference*, 1–3 December 1999, Anchorage, Alaska, 127–142.
- Schumacher, J.D., and V. Alexander (1999): Variability and the role of the physical environment in the Bering Sea ecosystem. In *Dynamics of the Bering Sea*, T.R. Loughlin, and K. Ohtani (eds.), University of Alaska Sea Grant, AK-SG-99-03, Fairbanks, Alaska, 147–160.
- Schumacher, J.D., N.A. Bond, R.D. Brodeur, P.A. Livingston, J.M. Napp, and P.J. Stabeno (2003): Climate change in the southeastern Bering Sea and some consequences for biota. In *Large Marine Ecosystems of the World: Trends in Exploitation, Protection and Research*, G. Hemplé, and K. Sherman (eds.), Amsterdam: Elsevier Science, 17–40.
- Schumacher, J.D., and T.H. Kinder (1983): Low frequency currents over the Bering Sea Shelf. *J. Phys. Oceanogr.*, 13, 607–623.
- Schumacher, J.D., and R.K. Reed (1992): Characteristics of currents over the continental slope of the eastern Bering Sea. *J. Geophys. Res.*, 97, 9423–9433.
- Schumacher, J.D., and P.J. Stabeno (1994): Ubiquitous eddies of the eastern Bering Sea and their coincidence with concentrations of larval pollock. *Fish. Oceanogr.*, 3, 182–190.
- Schumacher, J.D., and P.J. Stabeno (1998): Continental shelf of the Bering Sea. In *The Sea, Vol. 11, The Global Coastal Ocean: Regional Studies and Synthesis*, John Wiley & Sons, Inc., New York, NY, 789–822.
- Schwinger, F.B., and C.S. Moore (2000): A year without summer for California: Or a harbinger of a climate shift. *Eos Trans. AGU*, 81, 301.
- SEBSCC (1995): Concept Paper. <http://www.pmel.noaa.gov/sebscc>.
- Shchepetkin, A.F., and J.C. McWilliams (1998): Quasi-monotone advection schemes based on explicit locally adaptive dissipation. *Mon. Weather Rev.*, 126, 1541–1580.
- Sherr, E.B., and B.F. Sherr (1992): Trophic roles of pelagic protists: Phagotrophic flagellates as herbivores. *Arch. Hydrobiol. Beih. Ergebn. Limnol.*, 37, 165–172.
- Shuer, P.G., and J.J. Walsh (1993): A coupled physical-biological model of the Bering-Chukchi Seas. *Cont. Shelf Res.*, 13, 543–574.
- Sinclair, E.H., G.A. Antonelis, B.W. Robson, R.R. Ream, and T.R. Loughlin (1996): Northern fur seal, *Callorhinus ursinus*, predation on juvenile walleye pollock, *Theragra chalcogramma*. In *Ecology of Juvenile Walleye Pollock, Theragra chalcogramma*, R.D. Brodeur, P.A. Livingston, T.R. Loughlin, and A.B. Hollowed (eds.), NOAA Tech. Rept. NMFS 126, 167–178.
- Sinclair, E.H., A.A. Balanov, T. Kubodera, V.I. Radchenko, and Y.A. Fedorets (1999): Distribution and ecology of mesopelagic fishes and cephalopods. In *Dynamics of the Bering Sea*, T.R. Loughlin, and K. Ohtani (eds.), University of Alaska Sea Grant, AK-SG-99-03, Fairbanks, Alaska, 485–508.
- * Sinclair, E.H., and P.J. Stabeno (2002): Mesopelagic nekton and associated physics of the southeastern Bering Sea. *Deep-Sea Res. II, Top. Stud. Oceanogr.*, 49(26), 6127–6145.
- Sinclair, E.H., T. Loughlin, and W. Pearcy (1994): Prey selection by northern fur seals (*Callorhinus ursinus*) in the eastern Bering Sea. *Fish. Bull.*, 92, 144–156.
- * Sinclair, E.H., G.L. Hunt, Jr., L. Vlietstra, T.K. Zeppelin, G.V. Byrd, and A.M. Springer: Trends in diet and population of apex predators in the Pribilof Islands: 1975–2000. *Can. J. Fish. Oceanogr.*, in preparation.

- * Sinclair, E.H., T.K. Zeppelin, and R.R. Ream (2003): Synthesis of pinniped diets at the Pribilof Islands. Final Report of the Synthesis of Information on Marine Birds, Fur Seals and Steller Sea Lions in the Eastern Bering Sea. SEBSCC report, NOAA/PMEL, 7600 Sand Point Way NE, Seattle, WA 98115.
- * Smith, S.L., S.M. Henrichs, and T. Rho (2002): Stable C and N isotopic composition of sinking particles and zooplankton over the southeastern Bering Sea shelf. *Deep-Sea Res. II, Top. Stud. Oceanogr.*, 49(26), 6031–6050.
- Smith, S.L., and J. Vidal (1984): Spatial and temporal effects of salinity, temperature, and chlorophyll on the communities of zooplankton in the southeastern Bering Sea. *J. Mar. Res.*, 42, 221–257.
- Smith, S.L., and J. Vidal (1986): Variations in the distribution, abundance and development of copepods in the southeast Bering Sea in 1980 and 1981. *Cont. Shelf Res.*, 5, 215–239.
- Smith, S.L., S.M. Henrichs, and T. Rho (2002): Stable C and N isotopic composition of sinking particles and zooplankton over the southeastern Bering Sea shelf. *Deep-Sea Res. II, Top. Stud. Oceanogr.*, 49, 6031–6050.
- Sogard, S.M., and B.L. Olla (1993): Effects of light, thermoclines and predator presence on vertical distribution and behavioral interactions of juvenile walleye pollock, *Theragra chalcogramma* Pallas. *J. Exp. Mar. Biol. Ecol.*, 167, 179–195.
- Springer, A.M. (1991): Seabird relationships to food webs and the environment: examples from the North Pacific Ocean. In *Studies of High-Latitude Seabirds. 1. Behavioral, Energetic, and Oceanographic Aspects of Seabird Feeding Ecology*, W.A. Montevecchi, and A.J. Gaston (eds.), Ottawa, Canadian Wildlife Service, 39–48.
- Springer, A.M. (1992): A review: Walleye pollock in the North Pacific—how much difference do they really make? *Fish. Oceanogr.*, 1, 80–96.
- Springer, A.M. (1998): Is it all climate change? Why marine bird and mammal populations fluctuate in the North Pacific. In *Proceedings of the 10th 'Aha Huliko'a Hawaiian Winter Workshop on Biotic Impacts of Extratropical Climate Variability in the Pacific*, 26–30 January 1998, G. Holloway, P. Muller, and D. Henderson (eds.), SOEST Special Publication, University of Hawaii, Honolulu, HI, 121–125.
- Springer, A.M., and G.V. Byrd (1989): Seabird dependence on walleye pollock in the southeastern Bering Sea. In *Proceedings of the International Symposium on the Biology and Management of Walleye Pollock*, Alaska Sea Grant, AK-SG-89-01, 667–677.
- Springer, A.M., and C.P. McRoy (1993): The paradox of pelagic food webs in the northern Bering Sea—III. Patterns of primary production. *Cont. Shelf Res.*, 13, 575–599.
- Springer, A.M., and D.G. Roseneau (1985): Copepod-based food webs: Auklets and oceanography in the Bering Sea. *Mar. Ecol. Prog. Ser.*, 21, 229–237.
- Springer, A.M., C.P. McRoy, and M. Flint (1996a): The Bering Sea green belt: shelf-edge processes and ecosystem production. *Fish. Oceanogr.*, 5, 205–223.
- Springer, A.M., C.P. McRoy, and K.R. Turco (1989): The paradox of pelagic food webs in the northern Bering Sea—II. Zooplankton communities. *Cont. Shelf Res.*, 9, 359–386.
- Springer, A.M., E.C. Murphy, D.G. Roseneau, C.P. McRoy, and B.A. Cooper (1987): The paradox of food webs in the northern Bering Sea—I. Seabird food habits. *Cont. Shelf Res.*, 7(8), 895–911.
- Springer, A.M., J.F. Piatt, and G.B. Van Vliet (1996): Seabirds as proxies of marine habitats and food webs in the western Aleutian Arc. *Fish. Oceanogr.*, 5, 45–55.

*SEBSCC publication

- * Stabeno, P.J., N.A. Bond, N.B. Kachel, S.A. Salo, and J.D. Schumacher (2001): On the temporal variability of the physical environment over the south-eastern Bering Sea. *Fish. Oceanogr.*, 10(1), 81–98.
- * Stabeno, P.J., and G.L. Hunt, Jr. (2002): Overview of the Inner Front and South-east Bering Sea Carrying Capacity Programs. *Deep-Sea Res. II, Top. Stud. Oceanogr.*, 49(26), 6157–6168.
- * Stabeno, P.J., N.B. Kachel, M.E. Sullivan, and T.E. Whitledge (2002): Variability of physical and chemical characteristics along the 70-m isobath of the south-eastern Bering Sea. *Deep-Sea Res. II, Top. Stud. Oceanogr.*, 49(26), 5931–5943.
- * Stabeno, P.J., and J.E. Overland (2001): The Bering Sea shifts toward an earlier Spring transition. *Eos Trans. AGU*, 29, 317, 321.
- Stabeno, P.J., and R.K. Reed (1994): Circulation in the Bering Sea basin observed by satellite-tracked drifters: 1986–1993. *J. Phys. Oceanogr.*, 24, 848–854.
- * Stabeno, P.J., R.K. Reed, and J.M. Napp (2002): Transport through Unimak Pass, Alaska. *Deep-Sea Res. II, Top. Stud. Oceanogr.*, 49(26), 5919–5930.
- Stabeno, P.J., J.D. Schumacher, R.F. Davis, and J.M. Napp (1998): Under-ice observations of water column temperature, salinity and spring phytoplankton dynamics: Eastern Bering Sea shelf, 1995. *J. Mar. Res.*, 56, 239–255.
- Stabeno, P.J., J.D. Schumacher, and K. Ohtani (1999a): The physical oceanography of the Bering Sea. In *Dynamics of The Bering Sea*, T.R. Loughlin, and K. Ohtani (eds.), Alaska Sea Grant, AK-SG-99-03, Fairbanks, AK, 1–28.
- Stabeno, P.J., J.D. Schumacher, S.A. Salo, G.L. Hunt, Jr., and M. Flint (1999b): Physical environment around the Pribilof Islands. In *Dynamics of the Bering Sea*, T.R. Loughlin, and K. Ohtani (eds.), Alaska Sea Grant, AK-SG-99-03, Fairbanks, AK, 193–215.
- Stabeno, P.J., and P. Van Meurs (1999): Evidence of episodic on-shelf flow in the southeastern Bering Sea. *J. Geophys. Res.*, 104(C12), 29,715–29,720.
- * Stockwell, D.A., T.E. Whitledge, S.I. Zeeman, K.O. Coyle, J.M. Napp, R.D. Brodeur, A.I. Pinchuk, and G.L. Hunt, Jr. (2001): Anomalous conditions in the south-eastern Bering Sea, 1997: Nutrients, phytoplankton and zooplankton. *Fish. Oceanogr.*, 10(1), 99–116.
- Sugimoto, T., and K. Tadokoro (1997): Interannual-interdecadal variations in zooplankton biomass, chlorophyll concentration and physical environment in the subarctic Pacific and Bering Sea. *Fish. Oceanogr.*, 6, 74–93.
- Sugimoto, T., and K. Tadokoro (1998): Interdecadal variations of plankton biomass and physical environment in the North Pacific. *Fish. Oceanogr.*, 7, 289–299.
- Sukhanova, I.N., and M.V. Flint (1998): Anomalous blooming of cocolithophorids over the eastern Bering Sea shelf. *Oceanography*, 38, 502–505.
- Sukhanova, I.N., H.J. Semina, and M.V. Venttsel (1999): Spatial distribution and temporal variability of phytoplankton in the Bering Sea. In *Dynamics of the Bering Sea*, T.R. Loughlin, and K. Ohtani (eds.), University of Alaska Sea Grant, AK-SG-99-03, Fairbanks, AK, 193–215.
- Sverdrup, H.U. (1953): On conditions for the vernal blooming of phytoplankton. *J. Conseil Exp. Mer*, 18, 287–295.
- * Swartzman, G.L., R.D. Brodeur, J.M. Napp, G.L. Hunt, Jr., D. Demer, and R.P. Hewitt (1999): Spatial proximity of age-0 walleye pollock (*Theragra chalcogramma*) to zooplankton near the Pribilof Islands, Bering Sea, Alaska. *ICES J. Mar. Sci.*, 56(4), 545–560.
- * Swartzman, G.L., and G.L. Hunt, Jr. (2000): Spatial association between murres (*Uria* spp.), puffins (*Fratercula* spp.) and fish shoals near the Pribilof Islands, Alaska. *Mar. Ecol. Prog. Ser.*, 206, 297–309.

- * Swartzman, G.L., J.M. Napp, R.D. Brodeur, A.G. Winter, and L. Ciannelli (2002): Spatial patterns of pollock and zooplankton distribution in the Pribilof Islands, Alaska nursery area and their relationship to pollock recruitment. *ICES J. Mar. Sci.*, 59(6), 1167–1186.
- * Swartzman, G.L., A.G. Winter, K.O. Coyle, R.D. Brodeur, T. Buckley, L. Ciannelli, G.L. Hunt, Jr., J. Ianelli, and S.A. Macklin (2004): Relationship of age-0 pollock abundance and distribution around the Pribilof Islands with other shelf regions of the eastern Bering Sea. *Fish. Res.* (submitted).
- Thompson, D.W.J., and J.M. Wallace (1998): The Arctic Oscillation signature in the wintertime geopotential height and temperature fields. *Geophys. Res. Lett.*, 25, 1297–1300.
- Townsend, D.W., L.M. Cammen, P.M. Holligan, D.E. Campbell, and N.R. Pettigrew (1994a): Causes and consequences of variability in the timing of spring phytoplankton blooms. *Deep-Sea Res.*, 41, 747–765.
- Townsend, D.W., M.D. Keller, P.M. Holligan, S.G. Ackleson, and W.M. Balch (1994b): Blooms of the coccolithophore *Emiliania huxleyi* with respect to hydrography in the Gulf of Maine. *Cont. Shelf Res.*, 14, 979–1000.
- Traynor, J., and D. Smith, 1996. Summer distribution and abundance of age-0 walleye pollock, *Theragra chalcogramma*, in the Bering Sea. (Abstract). In *Ecology of Juvenile Walleye Pollock, Theragra chalcogramma*, NOAA Tech. Rept. NMFS 126.
- Trenberth, K.E. and J.W. Hurrell (1994): Decadal atmospheric-ocean variations in the North Pacific. *Clim. Dyn.*, 9, 303–319.
- Trenberth, K.E., and J.W. Hurrell (1995): Decadal coupled atmospheric-ocean variations in the North Pacific Ocean. In *Climate Change and Northern Fish Populations*, R.J. Beamish (ed.), *Can. Spec. Publ. Fish. Aquat. Sci.*, 121, 15–24.
- Trites, A.W., and M.A. Bigg (1996): Physical growth of northern fur seals: Seasonal fluctuations and migratory influences. *J. Zool. (Lond.)*, 238, 459–482.
- Trites, A.W., and P.A. Larkin (1996): Changes in the abundance of Steller sea lions (*Eumetopias jubatus*) in Alaska from 1956 to 1992: How many were there? *Aquat. Mamm.*, 22, 153–166.
- Tynan, C.T. (1998): Coherence between whale distributions, chlorophyll concentration, and oceanographic conditions on the southeast Bering Sea shelf during a coccolithophore bloom, July–August, 1997. *Eos Trans. AGU*, 79, 127.
- Tynan, C.T. (1999): Redistribution of cetaceans in the southeast Bering Sea relative to anomalous oceanographic conditions during the 1997 El Niño. In *Proceedings of the 1998 Science Board Symposium on the Impacts of the 1997–1998 El Niño Event on the North Pacific Ocean and Its Marginal Seas*, H.J. Freeland, W.T. Peterson, and A. Tyler (eds.), PICES Scientific Report 10, 115–117.
- USGS (United States Geological Survey), (2004): Investigations of the linkages between climate, growth, and production of sockeye salmon populations in Bristol Bay, Alaska. http://www.nrel.colostate.edu/brd_global_change/proj_25_ak_salmon.html. March 17, 2004.
- * Vance, T.C., C.T. Baier, R.D. Brodeur, K.O. Coyle, M.B. Decker, T. Wyllie-Echeverria, G.L. Hunt, Jr., J.M. Napp, J.D. Schumacher, P.J. Stabeno, D.A. Stockwell, C. Tynan, T.E. Whitledge, and S.I. Zeeman (1998): Union ecosystem anomalies in the eastern Bering Sea: Including blooms, birds and other biota. *Eos Trans. AGU*, 79(10), 121, 126.
- * Vance, T.C., and S.A. Macklin (1998): Anomalies of the eastern Bering Sea ecosystem during Summer and Fall 1997. *Alaska Fish. J.*, 21(3), 36–37.
- Vidal, J. (1980): Physioecology of zooplankton II: Effects of phytoplankton concen-

- tration, temperature, and body size on the development and molting rates of *Calanus pacificus* and *Pseudocalanus* spp. *Mar. Biol.*, 56, 135–146.
- Vidal, J., and S.L. Smith (1986): Biomass, growth and development of population of herbivorous zooplankton in the southeastern Bering Sea. *Deep-Sea Res.*, 33, 523–566.
- * Vlietstra, L., A. Springer, G.V. Byrd, G.L. Hunt, Jr. (2003). Control of seabird population dynamics at the Pribilof Islands, Alaska. Final Report of the Synthesis of Information on Marine Birds, Fur Seals and Steller Sea Lions in the Eastern Bering Sea. SEBSCC report, NOAA/PMEL, 7600 Sand Point Way NE, Seattle, WA 98115.
- Wallace, J.M., C. Smith, and C.S. Bretherton (1992): Singular value decomposition of wintertime sea surface temperature and 500-hPa height anomalies. *J. Clim.*, 5, 561–576.
- Walsh, J.J., and C.P. McRoy (1986): Ecosystem analysis in the southeastern Bering Sea. *Cont. Shelf Res.*, 5, 259–288.
- Welch, D.W., B.R. Ward, B.D. Smith, and J.P. Eveson (1998): Influence of the 1990 ocean climate shift on British Columbia steelhead (*O. mykiss*) and coho (*O. kisutch*) populations. In *Proceedings of the 10th 'Aha Huliko'a Hawaiian Winter Workshop on Biotic Impacts of Extratropical Climate Variability in the Pacific*, 26–30 January 1998, G. Holloway, P. Muller, and D. Henderson, (eds.), SOEST Special Publication, University of Hawaii, Honolulu, HI, 77–87.
- Wespestad, V.G. (1994): Walleye Pollock. Report and data submitted for the Third International Pollock Assessment Workshop, Seattle, WA, 119–143.
- Wespestad, V.G., L.W. Fritz, W.J. Ingraham, Jr., and B.A. Megrey (2000): On relationships between cannibalism, climate variability, physical transport and recruitment success of Bering Sea walleye pollock, *Theragra chalcogramma*. *ICES J. Mar. Sci.*, 57, 272–278.
- Wespestad, V.G., and T.J. Quinn, II (1996): Importance of cannibalism in the population dynamics of walleye pollock, *Theragra chalcogramma*. In *Ecology of Juvenile Walleye Pollock Theragra chalcogramma*, R.D. Brodeur, P.A. Livingston, T.R. Loughlin, and A.B. Hollowed (eds.), NOAA Tech. Rept. NMFS 126, 212–226.
- * Whittlestone, T.E., and V.A. Luchin (1999): Summary of chemical distributions and dynamics in the Bering Sea. In *Dynamics of the Bering Sea*, T.R. Loughlin, and K. Ohtani (eds.), University of Alaska Sea Grant, Fairbanks, Alaska, AK-SG-99-03, 217–250.
- * Wilderbuer, T.K., A.B. Hollowed, W.J. Ingraham, Jr., P.D. Spencer, M.E. Connors, N.A. Bond, and G.E. Walters (2002): Flatfish recruitment response to decadal climatic variability and ocean conditions in the eastern Bering Sea. *Prog. Oceanogr.*, 55(1–2), 235–247.
- * Wilson, M.T., A.H. Andrews, A.L. Brown, and E.E. Cordes (2002): Axial rod growth and age estimation of the sea pen, *Halipteris willemoesi köllikeri*. *Hydrobiologia*, 471(1–3), 133–142.
- * Winter, A.G., G.L. Swartzman, and L. Ciannelli. Early- to late-summer population growth and prey consumption by age-0 pollock in two years of contrasting pollock abundance near the Pribilof Islands, Bering Sea (submitted).
- Wyllie-Echeverria, T. (1995): Sea-ice conditions and the distribution of walleye pollock (*Theragra chalcogramma*) on the Bering and Chukchi shelf. In *Climate Change and Northern Fish Populations*, R.J. Beamish, (ed.), *Can. Spec. Publ. Fish. Aquat. Sci.*, 121, 131–136.
- Wyllie-Echeverria, T. (1996): The relationship between the distribution of one-year-old walleye pollock, *Theragra chalcogramma*, and sea-ice characteristics.

- In *Ecology of Juvenile Walleye Pollock, Theragra chalcogramma*, R.D. Brodeur, P.A. Livingston, T.R. Loughlin, and A.B. Hollowed (eds.), NOAA Tech. Rept. NMFS 126, 47–56.
- Wyllie-Echeverria, T., and K. Ohtani (1999): Seasonal sea ice variability and the Bering Sea ecosystem. In *Dynamics of the Bering Sea*, T.R. Loughlin, and K. Ohtani (eds.), University of Alaska Sea Grant, AK-SG-99-03, Fairbanks, AK, 435–451.
- Wyllie-Echeverria, T., and W.S. Wooster (1998): Year-to-year variations in Bering Sea ice cover and some consequences for fish distributions. *Fish. Oceanogr.*, 7, 159–170.
- York, A.E., and J.R. Hartley (1981): Pup production following harvest of female northern fur seals. *Can. J. Fish. Aquat. Sci.*, 38, 84–90.
- York, A.E., and P. Kozloff (1987): On the estimation of numbers of northern fur seal, *Callorhinus ursinus*, pups born on St. Paul Island, 1980–86. *Fish. Bull.*, 85, 367–375.
- York, A.E., R.G. Towell, R.R. Ream, J.D. Baker, and B.W. Robson (2000): Population assessment, Pribilof Islands, Alaska. In *Fur Seal Investigations, 1998*, B.W. Robson, (ed.), NOAA Tech. Memo. NMFS-AFSC-113, 7–26.